

Spontaneous localization of bulk matter fields

S. Groot Nibbelink ^{a,1}, H.P. Nilles ^{a,2}, M. Olechowski ^{a,b,3}

^a *Physikalisches Institut der Universität Bonn,
Nussallee 12, 53115 Bonn, Germany.*

^b *Institute of Theoretical Physics, Warsaw University,
Hoża 69, 00-681 Warsaw, Poland.*

Abstract

We study models compactified on $S^1/\mathbb{Z}_2$ with bulk and brane matter fields charged under $U(1)$ gauge symmetry. We calculate the FI-terms and show by minimizing the resulting potential that supersymmetry or gauge symmetry is spontaneously broken if the sum of the charges does not vanish. Even if this sum vanishes, there could be an instability as a consequence of localized FI-terms. This leads to a spontaneous localization of charged bulk fields on respective branes.

¹ E-mail: nibblink@th.physik.uni-bonn.de

² E-mail: nilles@th.physik.uni-bonn.de

³ E-mail: olech@th.physik.uni-bonn.de

Theories in more than 4-dimensional space-time have attracted much attention in recent years. The connection to our 4-dimensional world is made through compactification of extra space dimensions. Among the proposed mechanisms, orbifold compactification seems to be particularly simple and instructive [1]. In addition, it allows the incorporation of so-called brane world scenarios, in which branes are placed (at fixed points) in a higher dimensional bulk. We can then distinguish between bulk-fields that propagate in the higher dimensional bulk and brane fields whose propagation and interactions are confined to the respective branes (fixed points).

Various models of this kind have been proposed recently because of their potential phenomenological merits. Some unwanted couplings can be made very small or even zero by appropriate localization of the fields. For example, two fields localized at different branes can not interact directly but only indirectly via bulk fields. In addition, two such brane fields could have different couplings to a bulk field if that field has nontrivial profile along the extra dimensions. This type of mechanism could be responsible for the suppression of proton decay, Yukawa couplings and supersymmetry breakdown.

Many of these models assume some amount of supersymmetry to render the theory ultraviolet (UV) insensitive and solve the hierarchy problem. While in 4 dimensions the UV-behavior of the theory is well understood there remain some open questions in the brane world picture. It is one of these questions that we shall study in the present paper: the UV-sensitivity of the low-energy physics.

In 4 dimensions the supersymmetric models (also with soft breaking interactions) are usually at most logarithmically sensitive to the high scale Λ . The only exception (the Achilles' heel of supersymmetry) is the quadratically divergent Fayet-Iliopoulos term for a $U(1)$ gauge symmetry if the sum over the scalar fields of the corresponding charge is nonzero [2]. Thus, in phenomenologically acceptable models the $U(1)$ charges must sum up to zero; otherwise the low energy physics will be destabilized and either gauge symmetry or supersymmetry will be broken at a scale of order Λ . Another (independent) reason for this condition is the absence of the mixed gauge-gravitational anomaly if gravity is taken into account.

The Standard Model of strong and electroweak interactions contains a $U(1)$ gauge symmetry – hypercharge symmetry. The sum of hypercharges vanishes for each generation of fermions so also for each generation of scalars in the supersymmetric version of the model. Thus, there are no quadratically divergent FI-terms in the 4-dimensional supersymmetric extension of the standard model.

We would now like to ask the question how this situation generalizes to the higher dimensional brane world scenario. Previously [3] we have already shown that the presence of massless bulk scalar fields with nonvanishing sum of charges leads to a quadratic divergence⁴. In this paper we set up the general case with

⁴For a related discussion of ultraviolet sensitivity in extra dimensions see [4, 5, 6].

charged bulk and brane fields and arrive at a similar conclusion: if the sum of charges does not add up to zero, there is a quadratically divergent FI-term and either supersymmetry or gauge symmetry are destabilized in the same way as in the 4-dimensional theory.

Still this is not the full story. We next concentrate on the more specific question of localization of the field with nonzero hypercharge. Usually in brane world models some of the standard model particles can be described by bulk fields, while others by brane fields. Thus generically the hypercharges add up to zero only **globally** (when integrated over the extra dimension) but not **locally**. In such a situation we can show that as a result of the localized FI-terms we find a **spontaneous localization** of charged bulk fields. The zero modes of these bulk fields become localized on the brane, while the masses of the higher Kaluza-Klein modes are pushed to the UV-scale Λ .

In the present paper we shall present our results in the framework of the simplest possible toy model. The general results for more realistic models will be presented in a future publication [7]. We consider the 5-dimensional supersymmetric model compactified on $S^1/\mathbb{Z}_2$ containing 3 multiplets: two 5-dimensional bulk multiplets – the vector multiplet V and the hypermultiplet H – and one 4-dimensional chiral multiplet C_0 localized at the brane at $y = 0$. One could expect a gauge anomaly in such a model similar to the one discussed in [8]. However we will postpone the discussion of the gauge anomalies here for two reasons. First, the simple model allows a simplest possible illustration of the mechanism of localization observed here, and secondly, the localization of fields might actually lead to a more refined and subtle discussion of gauge anomalies, including a possible cancellation mechanism via Chern-Simons terms [9].

The 5-dimensional off-shell vector multiplet $V = (A_M, \lambda^i, \Phi, \vec{D})$ contains a gauge field A_M , a doublet of symplectic-Majorana gauginos λ^i , a real scalar Φ and a triplet of auxiliary fields $\vec{D}$ (we use the standard notation: $M = 1 \dots 5$, $\mu = 1 \dots 4$, $x^5 = y$). Each of the components must have a definite parity under the $\mathbb{Z}_2$ symmetry used for orbifolding. We choose the parity assignments as follows:

$$V : \begin{array}{|c|c|c|c|c|c|c|} \hline \text{state} & A_\mu & A_5 & \Phi & \lambda_\pm & D^3 & D^{1,2} \\ \hline \text{parity} & + & - & - & \pm & + & - \\ \hline \end{array}$$

With this choice neither the $N = 1$ supersymmetry nor the $U(1)$ gauge symmetry is broken by orbifolding.

The 5-dimensional off-shell hypermultiplet H consists of two complex scalars $\phi_\pm$, one Dirac fermion ψ , and two complex auxiliary fields $F_\pm$. The subscripts on the bosonic components denote their parity under $\mathbb{Z}_2$: $\phi_\pm(-y) = \pm\phi_\pm(y)$, $F_\pm(-y) = \pm F_\pm(y)$. The parity of the components of ψ is related to their 4-dimensional chirality: $\psi(-y) = i\gamma^5\psi(y)$.

The bulk interactions of V and H are described by the standard 5-dimensional supersymmetric Lagrangian. The part of that Lagrangian which is important for

our discussion can be written, in terms of the above introduced fields, as:

$$\begin{aligned}\mathcal{L}_{\text{bulk}} = & -\frac{1}{2}(\partial_M\Phi)^2 + \frac{1}{2}\vec{D}^2 - \mathcal{D}_M\phi_+^\dagger\mathcal{D}^M\phi_+ - \mathcal{D}_M\phi_-^\dagger\mathcal{D}^M\phi_- + i\bar{\psi}\gamma^M\mathcal{D}_M\psi \\ & - gqD^3\left(\phi_+^\dagger\phi_+ - \phi_-^\dagger\phi_-\right) - gq\left((D^1 - iD^2)\phi_+^T\phi_- + \text{h.c.}\right) \\ & - g^2q^2\Phi^2\left(\phi_+^\dagger\phi_+ + \phi_-^\dagger\phi_-\right) - gq\Phi\bar{\psi}\psi + \dots\end{aligned}\quad (1)$$

Now we add a 4-dimensional chiral multiplet $C_0 = (\phi_0, \psi_0, F_0)$ localized at the brane at $y = 0$. It was shown in ref. [10] that such a brane multiplet can be coupled to the bulk gauge multiplet in a way which preserves $N = 1$ supersymmetry. The coupling is given by the standard 4-dimensional interaction of a chiral multiplet with a gauge multiplet when the role of a 4-dimensional gauge multiplet is played by the boundary values of the appropriate components of the gauge bulk field⁵: $(A_\mu, \lambda_+, D^3 - \partial_y\Phi)$. The resulting Lagrangian reads

$$\mathcal{L}_{\text{brane}} = \delta(y)\left(-\mathcal{D}_\mu\phi_0^\dagger\mathcal{D}^\mu\phi_0 + gq_0(D^3 - \partial_y\Phi)\phi_0^\dagger\phi_0 + \dots\right)\quad (2)$$

where q_0 is the $U(1)$ charge of C_0 . The tree level action of the model is given by the sum of the bulk and brane contributions

$$S^{(0)} = \int d^4x dy (\mathcal{L}_{\text{bulk}} + \mathcal{L}_{\text{brane}})\quad (3)$$

and is invariant under the $N = 1$ supersymmetry and the $U(1)$ gauge symmetry.

Let us now take the 1-loop corrections into account. It occurs that divergent FI-terms play a very crucial role. They are generated by tadpole diagrams even if the sum of all charges is zero because the charge is non-trivially distributed along the fifth dimension.

We start with the contribution coming from the tadpole with the brane scalar ϕ_0 in the loop (fig. 1a). It is given by the standard 4-dimensional result with the gauge auxiliary field replaced by the boundary value of the combination $(D^3 - \partial_y\Phi)$. In the cut-off regularization it is given by

$$(D^3 - \partial_y\Phi) gq_0 \frac{\Lambda^2}{16\pi^2} \delta(y).\quad (4)$$

The calculation of the bulk field contributions is more involved (part of it has been already presented in refs. [3, 8]). It can be performed by expanding the bulk fields into 4-dimensional modes and summing the contributions from the obtained tadpole diagrams. The two kinds of bulk tadpole diagrams are

⁵ In a similar way the brane chiral multiplet can be coupled to the chiral multiplet made out of the boundary values of the bulk hypermultiplet components $(\phi_+, \psi_+, F_+ - \partial_y\phi_-)$. More details will be presented in a future publication [7].

shown on figs. 1b and 1c. In one of them the D^3 field is coupled to the scalar loops and only this contribution has been taken into account in the previous calculations [3, 8]. But there is also a contribution in which the Φ scalar is coupled to the fermion loop. It gives the same FI-term for $(-\partial_y \Phi)$ as the bulk scalar tadpole gives for D^3 . One should expect this because the model is invariant under $N = 1$ supersymmetry and the combination $(D^3 - \partial_y \Phi)$ is the gauge auxiliary field under this supersymmetry [10]. The details of the calculation will be presented elsewhere [7] and here we give only the final result:

$$(D^3 - \partial_y \Phi) g \frac{q}{2} \left\{ \frac{\Lambda^2}{16\pi^2} [\delta(y) + \delta(y - \pi R)] + \frac{\ln \Lambda^2}{16\pi^2} [\delta''(y) + \delta''(y - \pi R)] + \dots \right\} \quad (5)$$

where dots denote the finite contribution which will not be discussed here because we are interested only in the sensitivity of the model to the high energy scale represented by the cut-off Λ .

Let us now analyse our model with those 1-loop corrections quadratically and logarithmically sensitive to the cut-off scale Λ . We start with the 4-dimensional effective potential which is given by the sum of the potential part of the tree level action (3) and the integrals of (4) and (5). The part of the potential important for our analysis (i.e. without the F -type terms) reads

$$\begin{aligned} V = \int dy \left[-\frac{1}{2} \vec{D}^2 + \frac{1}{2} (\partial_y \Phi)^2 + |\partial_y \phi_+|^2 + |\partial_y \phi_-|^2 + \xi (D^3 - \partial_y \Phi) \right. \\ \left. + gq_0 (D^3 - \partial_y \Phi) |\phi_0|^2 \delta(y) + g^2 q^2 \Phi^2 (|\phi_+|^2 + |\phi_-|^2) \right. \\ \left. + gq D^3 (|\phi_+|^2 - |\phi_-|^2) + gq ((D^1 - iD^2) \phi_+ \phi_- + \text{h.c.}) \right] \quad (6) \end{aligned}$$

where the divergent radiatively generated FI-parameter $\xi(y)$ is given by

$$\xi = g \frac{\Lambda^2}{16\pi^2} \left[\left(q_0 + \frac{q}{2} \right) \delta(y) + \frac{q}{2} \delta(y - \pi R) \right] + g \frac{\ln \Lambda^2}{16\pi^2} \left[\frac{q}{2} \delta''(y) + \frac{q}{2} \delta''(y - \pi R) \right]. \quad (7)$$

We have dropped the A_5 component of the gauge field: as A_5 vanishes at both boundaries, we can perform a gauge transformation to put it zero over the bulk.

We rewrite this potential in a form more suitable for our later discussion. We add and subtract the term $gq(\partial_y \Phi) (|\phi_+|^2 - |\phi_-|^2)$ and perform an integration by parts, to make it possible to write the potential as a combination of squares:

$$\begin{aligned} V = \int dy \left[-\frac{1}{2} (-D^3 + \xi + gq_0 |\phi_0|^2 \delta(y) + gq |\phi_+|^2 - gq |\phi_-|^2)^2 \right. \\ \left. + \frac{1}{2} (-\partial_y \Phi + \xi + gq_0 |\phi_0|^2 \delta(y) + gq |\phi_+|^2 - gq |\phi_-|^2)^2 \right. \\ \left. - \frac{1}{2} |D^1 + iD^2 + gq \phi_+ \phi_-|^2 + g^2 q^2 |\phi_+ \phi_-|^2 \right. \\ \left. + |\partial_y \phi_+ - gq \Phi \phi_+|^2 + |\partial_y \phi_- + gq \Phi \phi_-|^2 \right]. \quad (8) \end{aligned}$$




Figure 1: The diagrams that give rise to FI-terms via brane scalar (a), bulk scalar (b) and bulk fermion (c) loops.

By splitting of the squares, it seems that we have introduced squares of delta functions or their derivatives. However, because of the opposite sign of the auxiliary field D^3 all this cancels out precisely. Thus the dangerous squares of $\delta(y)$ and $\delta''(y)$ do not appear. (Here we presented only this simplified argumentation and we postpone a more complete calculation to a future publication [7]). As the only negative terms in this potential are due to the auxiliary fields, that have algebraic equations of motion, for example

$$D^3(y) = \xi(y) + gq_0 |\phi_0|^2 \delta(y) + gq |\phi_+(y)|^2 - gq |\phi_-(y)|^2, \quad (9)$$

the potential (8) is clearly positive semi-definite.

Next, we investigate background field configurations, i.e. Vacuum Expectation Values (VEVs) of the scalar fields (which may be functions of y) that minimize the potential V . An important question is whether supersymmetry or $U(1)$ gauge symmetry is spontaneously broken. Gauge symmetry remains unbroken if the charged fields have vanishing VEVs: $\langle \phi_+ \rangle = \langle \phi_- \rangle = \langle \phi_0 \rangle = 0$, supersymmetry is unbroken if the potential vanishes in the minimum. As can be seen from the potential (8) this implies that

$$\langle D^3(y) \rangle - \partial_y \langle \Phi(y) \rangle = 0 \quad (10)$$

for a supersymmetric background field configuration. As this is a first order differential equation for the odd $\langle \Phi \rangle$, which vanishes at the boundaries, a supersymmetric VEV is only possible if the constant mode $D_0^3 = \int dy D^3(y)$ of the auxiliary field (9)

$$D_0^3 = \frac{g}{\pi R} \left((q + q_0) \frac{\Lambda^2}{16\pi^2} + q_0 |\langle \phi_0 \rangle|^2 + q \int_0^{\pi R} dy (|\langle \phi_+(y) \rangle|^2 - |\langle \phi_-(y) \rangle|^2) \right) \quad (11)$$

vanishes. (The logarithmic divergence is absent here, since the derivative $\delta'(y)$ vanishes at $y = 0$.) Hence, as far as the auxiliary field D_3 is concerned, only its zero mode D_0^3 is important for determining whether supersymmetric minimum can exist, in the presence of FI-terms localized at the branes in the 5-dimensional theory.

Let us consider two cases: *i*) the charges of the matter fields do not sum up to zero, $q + q_0 \neq 0$; *ii*) the charges sum up to zero, $q + q_0 = 0$. These cases are different because the radiatively generated quadratically divergent contribution to D_0^3 is proportional to the sum of charges $q + q_0$ (see (11)).

In case *i*) the supersymmetry can remain unbroken only when some nonzero VEVs of the charged fields cancel that Λ^2 contribution to D_0^3 :

$$(q + q_0) \frac{\Lambda^2}{16\pi^2} + q_0 |\langle \phi_0 \rangle|^2 + q \int_0^{\pi R} dy (|\langle \phi_+(y) \rangle|^2 - |\langle \phi_-(y) \rangle|^2) = 0. \quad (12)$$

The bulk field ϕ_0 can be used for this purpose if its charge q_0 has opposite sign with respect to the sum $q + q_0$ (i.e. $qq_0 < 0$ and $|q_0| < |q|$).

The situation with the bulk field is more complicated. We have to remember that its VEV must minimize the whole potential V and not only its first term depending on D^3 . Only one of the components, ϕ_+ or ϕ_- , can have nonvanishing VEV because of the $|\phi_+\phi_-|^2$ term in the potential (8). From the last two terms in V we see that such nonzero VEVs must satisfy

$$\partial_y \langle \phi_{\pm}(y) \rangle = \mp gq \langle \Phi(y) \rangle \langle \phi_{\pm}(y) \rangle. \quad (13)$$

In the case of $\langle \phi_+ \rangle$ it can be solved with the normalization chosen to fulfill eq. (12). The configuration with such $\langle \phi_+ \rangle$ and $\langle \phi_- \rangle = 0 = \langle \phi_0 \rangle$ gives vanishing value of V which is a global minimum because V is positive semi-definite. Nonzero $\langle \phi_+ \rangle$ can be used to restore supersymmetry if q has opposite sign with respect to the sum $q + q_0$ ($qq_0 < 0$ and $|q| < |q_0|$).

Equation (13) has no nonzero solutions for the odd field ϕ_- . The reason is that for $\lim_{y \rightarrow 0} \phi_-(y) \neq 0$ the l.h.s. of (13) has a $\delta(y)$ type singularity while no such structure appears at the r.h.s. of that equation. As the result $\langle \phi_- \rangle$ vanishes and can not cancel the Λ^2 term in D_0^3 .

We see that there are two possibilities in the case *i*) ($q + q_0 \neq 0$). If the bulk and brane field charges have opposite signs then one of these fields (the one with the smaller absolute value of the charge) develops a nonzero VEV. This VEV is such that the vacuum energy vanishes and the supersymmetry remains unbroken. But the $U(1)$ gauge symmetry is broken and the scale of breaking is given by the cut off scale Λ . If q and q_0 have the same sign then $U(1)$ is unbroken because the charged fields do not develop nonzero VEVs. (As argued above, the oppositely charged odd field ϕ_- cannot have a VEV.) In such a case the value of the vacuum energy is positive and supersymmetry is broken at very high scale given by Λ . Thus the situation is very similar to the 4-dimensional theories with quadratically divergent Fayet–Iliopoulos terms generated radiatively if the $U(1)$ charges do not sum up to zero.

Let us now go to the case *ii*) in which charges do sum up to zero: $q + q_0 = 0$. Now there is no Λ^2 contribution to D_0^3 given by (11). It is obvious that the potential V is minimized for vanishing VEVs of all matter fields. The gauge

$U(1)$ symmetry and the supersymmetry are unbroken. This is again very similar to the 4-dimensional theory but there are important differences: In 4 dimensions no FI-term is generated in such a case. In our bulk-brane setup nontrivial FI-like term is generated for the combination $(D^3 - \partial_y \Phi)$ and only its integral over y vanishes. The supersymmetry remains unbroken because the contributions to the VEV of the auxiliary field coming from D^3 and $\partial_y \Phi$ exactly cancel each other (this is related to the fact that Φ is a propagating field while D^3 is not). But we will see that the non-trivial FI-term for the propagating field Φ has very interesting consequences.

First we calculate the background configuration of Φ which is induced by the FI-terms. Using equations (7), (8) and (9) we get for $q_0 = -q$:

$$\langle \Phi(y) \rangle = -gq \frac{\Lambda^2}{64\pi^2} \epsilon(y) + gq \frac{\ln \Lambda^2}{32\pi^2} (\delta'(y) + \delta'(y - \pi R)) \quad (14)$$

where we again drop terms which are not sensitive to the scale Λ . Substituting this back to the part of the potential (8)

$$\int dy (|\partial_y \phi_+ - gq \langle \Phi \rangle \phi_+|^2 + |\partial_y \phi_- + gq \langle \Phi \rangle \phi_-|^2) \quad (15)$$

gives the mass terms for the fields ϕ_+ and ϕ_- in the effective 4-dimensional theory. The theory is supersymmetric so the nonzero VEV of Φ influences also the effective mass term for the matter fermions (via the last term in (1))

$$\int dy \bar{\psi} (i\gamma^5 \partial_y + gq \langle \Phi \rangle) \psi. \quad (16)$$

With vanishing $\langle \Phi \rangle$ the spectrum of $\phi_{\pm}$ is very simple: The even component has a constant massless mode while modes $\phi_+(y) = \cos(ny/R)$, $\phi_-(y) = \sin(ny/R)$ have masses equal to n/R . Let us now analyse the spectrum of the charged bulk fields with VEV of Φ given by (14). First of all the even field ϕ_+ has again a zero mode given formally by the formula

$$\phi_{+0}(y) = \exp \left(gq \int_0^y dy' \langle \Phi(y') \rangle \right) \phi_{+0}(0). \quad (17)$$

We can not directly substitute here $\langle \Phi \rangle$ as given by (14) because we would get arbitrary powers of $\delta'(y)$ which have no well definite meaning. We regularize the delta function with the appropriate Gaussian function so that

$$\delta'(y) = - \lim_{\sigma \rightarrow \infty} 2\pi\sigma^3 y \exp(-\pi\sigma^2 y^2). \quad (18)$$

Now the integral (17) can be calculated

$$\phi_{+0}(y) = N \exp \left\{ -\frac{g^2 q^2}{64\pi^2} \Lambda^2 y \right\} \exp \left\{ -\frac{g^2 q^2}{32\pi^2} \ln(\Lambda) \sigma \left[1 + e^{-\pi^2 \sigma^2} - e^{-\pi \sigma^2 y^2} - e^{-\pi \sigma^2 (y - \pi R)^2} \right] \right\}. \quad (19)$$

N is a normalization factor depending on Λ and σ . The expression in the second curly bracket is negative for $0 < y < \pi R$ and vanishes only at the fixed points. Thus the second exponential factor suppresses the zero mode very strongly for all points which are not close to $y = 0$ or $y = \pi R$. The first exponential factor involving Λ^2 suppresses further the zero mode for y away from 0. As a result the zero mode is exponentially localized at the $y = 0$ brane and the width of this localization is determined by the bigger of the scales Λ and σ . The cut off Λ can be treated as a scale characteristic for some more fundamental underlying theory, e.g. the string theory. The scale σ was introduced in order to resolve the infinitely thin branes. The thickness of the branes is likely to be determined by the scale characteristic for a theory describing brane dynamics (it could be related to Λ in a more fundamental theory). Anyway, the localization of the zero mode of ϕ_+ is not weaker than the localization of the brane itself (parameterized by σ). The FI-terms cause the zero mode ϕ_{+0} to become effectively a brane field localized at the same brane at which the oppositely charged field ϕ_0 is localized. Of course the corresponding fermionic zero mode of ψ is also localized at the same brane. This has important implications for the anomaly analysis in such models. For the case $q_0 = -q$ the anomaly is canceled locally at this brane (and is absent at the other brane).

We will not discuss in much detail the massive modes of the bulk fields. The important point is that they effectively disappear from the spectrum because their masses are at least as big as the cut off scale Λ . This can be seen by analyzing first the spectrum without the $\ln \Lambda$ term in (14). In such a case the mass eigenstates are given by

$$\begin{pmatrix} \phi_{+n} \\ \phi_{-n} \end{pmatrix} = \sqrt{\frac{2}{\pi R}} \begin{pmatrix} \cos [ny/R - \alpha] \\ \sin [ny/R] \end{pmatrix} \quad (20)$$

where the phase α is given by $\tan \alpha = gqR\Lambda^2/64\pi^2n$ and the corresponding masses are $m_n^2 = (gqR\Lambda^2/64\pi^2)^2 + (n/R)^2$. Then one can show that adding to $\langle \Phi \rangle$ a term depending on σ changes the profiles of the eigenstates but do not change their masses (because this new term integrates to zero and thus do not change the boundary conditions which are used to get the discrete spectrum).

A similar localization of the bulk fields takes place also when $q + q_0 \neq 0$. However the formulae in such a case are more complicated and will be discussed elsewhere [7].

We showed that the bulk matter field get localized at the brane due to the divergent FI-terms. One could ask the question whether such terms cause also the bulk gauge field to localize? The answer is: no. The gauge field have no self-couplings (it is an Abelian gauge symmetry) hence VEV of Φ does not change the ordinary Kaluza–Klein tower of states coming from the bulk gauge multiplet.

In the present paper we have analyzed the 5-dimensional model compactified on $S^1/\mathbb{Z}_2$ with one brane and one bulk matter multiplets coupled to the bulk

$U(1)$ gauge multiplet. It occurs that the Fayet–Iliopoulos terms are generated not only for the auxiliary component of the gauge multiplet, D^3 , but also for the derivative of the scalar component, $\partial_y \Phi$. The potential with these FI–terms added has been analyzed in order to check whether supersymmetry or gauge symmetry is spontaneously broken. The situation is very similar to the 4–dimensional case. Both supersymmetry and gauge symmetry remains unbroken only if the charges of matter fields sum up to zero. If the charges do not sum up to zero one of these symmetries is broken. Supersymmetry is broken if all the charges have the same sign. If charges of both signs exist then one of the matter fields develops a nonzero VEV and the gauge symmetry is broken. The scale of the breaking in both cases is of the order of the cut off scale Λ which should be regarded as a scale characteristic for an underlying more fundamental theory.

As a main result of this paper we have shown that the FI–terms have another very interesting consequences for this bulk–brane model. Namely, they lead to the spontaneous localization of the charged bulk matter field. The zero mode of the bulk field is localized at the brane at which the bulk field lives (if their charges have opposite signs). The width of such localization is related to the thickness of the brane. The massive modes of the bulk field get masses at least of the order of the cut off scale and disappear effectively from the spectrum. Thus, the bulk matter field changes spontaneously into a brane field. It seems that the model with the single bulk charged matter field develops this kind of instability.

We have discussed the simplest model with just one brane matter field, one bulk matter field and one bulk gauge field. However, the main features are very similar for more complicated models in which the $U(1)$ symmetry is just the hypercharge part of the Standard Model gauge group and there are many bulk and brane matter fields. We will consider more general models and their phenomenological applications in a future publication [7]. There we will also discuss in more detail consequences of this localization for the anomaly analysis.

Acknowledgments

Work supported in part by the European Community’s Human Potential Programme under contracts HPRN–CT–2000–00131 Quantum Spacetime, HPRN–CT–2000–00148 Physics Across the Present Energy Frontier and HPRN–CT–2000–00152 Supersymmetry and the Early Universe. SGN was supported by priority grant 1096 of the Deutsche Forschungsgemeinschaft. MO was partially supported by the Polish KBN grant 2 P03B 052 16.

References

- [1] L. J. Dixon, J. A. Harvey, C. Vafa and E. Witten, Nucl. Phys. B **261** (1985) 678.
- [2] W. Fischler, H. P. Nilles, J. Polchinski, S. Raby and L. Susskind, Phys. Rev. Lett. **47** (1981) 757.
- [3] D.M. Ghilencea, S. Groot Nibbelink and H.P. Nilles, Nucl. Phys. B **619** (2001) 385 [arXiv:hep-th/0108184].
- [4] D. M. Ghilencea and H. P. Nilles, Phys. Lett. B **507** (2001) 327 [arXiv:hep-ph/0103151].
- [5] S. Groot Nibbelink, Nucl. Phys. B **619** (2001) 373 [arXiv:hep-th/0108185].
- [6] D. M. Ghilencea, H. P. Nilles and S. Stieberger, arXiv:hep-th/0108183.
- [7] S. Groot Nibbelink, H.P. Nilles and M. Olechowski, to appear.
- [8] C.A. Scrucca, M. Serone, L. Silvestrini and F. Zwirner, Phys. Lett. B **525** (2002) 169 [arXiv:hep-th/0110073].
- [9] L. Pilo and A. Riotto, arXiv:hep-th/0202144.
- [10] E.A. Mirabelli and M.E. Peskin, Phys. Rev. D **58** (1998) 065002 [arXiv:hep-th/9712214].